




Contents lists available at ScienceDirect

Developmental Biology

journal homepage: www.elsevier.com/developmentalbiology

miR-17 family of microRNAs controls FGF10-mediated embryonic lung epithelial branching morphogenesis through MAPK14 and STAT3 regulation of E-Cadherin distribution

Gianni Carraro^a, Ahmed El-Hashash^a, Diego Guidolin^b, Caterina Tiozzo^a, Gianluca Turcatel^a, Brittany M. Young^a, Stijn P. De Langhe^a, Saverio Bellusci^a, Wei Shi^a, Pier Paolo Parnigotto^c, David Warburton^{a,*}

^a Developmental Biology, Regenerative Medicine and Surgery Program, Saban Research Institute, Children's Hospital Los Angeles, Keck School of Medicine and School of Dentistry, 4661 Sunset Boulevard MS 35, Los Angeles, CA 90027, USA

^b Department of Human Anatomy and Physiology, University of Padua, Padua, Italy

^c Department of Pharmaceutical Sciences, University of Padua, Padua, Italy

ARTICLE INFO

Article history:

Received for publication 13 April 2009

Revised 14 May 2009

Accepted 17 June 2009

Available online 25 June 2009

Keywords:

Lung
miRNAs
E-Cadherin
Structural homeostasis

ABSTRACT

The miR-17 family of microRNAs has recently been recognized for its importance during lung development. The transgenic overexpression of the entire miR-17–92 cluster in the lung epithelium led to elevated cellular proliferation and inhibition of differentiation, while targeted deletion of miR-17–92 and miR-106b–25 clusters showed embryonic or early post-natal lethality. Herein we demonstrate that miR-17 and its paralogs, miR-20a, and miR-106b, are highly expressed during the pseudoglandular stage and identify their critical functional role during embryonic lung development. Simultaneous downregulation of these three miRNAs in explants of isolated lung epithelium altered FGF10 induced budding morphogenesis, an effect that was rescued by synthetic miR-17. E-Cadherin levels were reduced, and its distribution was altered by miR-17, miR-20a and miR-106b downregulation, while conversely, beta-catenin activity was augmented, and expression of its downstream targets, including *Bmp4* as well as *Fgf2b*, increased. Finally, we identified *Stat3* and *Mapk14* as key direct targets of miR-17, miR-20a, and miR-106b and showed that simultaneous overexpression of *Stat3* and *Mapk14* mimics the alteration of E-Cadherin distribution observed after miR-17, miR-20a, and miR-106b downregulation. We conclude that the miR-17 family of miRNA modulates FGF10–FGFR2b downstream signaling by specifically targeting *Stat3* and *Mapk14*, hence regulating E-Cadherin expression, which in turn modulates epithelial bud morphogenesis in response to FGF10 signaling.

© 2009 Elsevier Inc. All rights reserved.

Introduction

Lung development is under the control of a complex network of regulatory proteins including those that participate in the FGF, TGFβ, SHH, and WNT pathways (Cardoso and Lu, 2006; Warburton et al., 2005). Branching and growth of the epithelium are principally controlled by the FGF10–FGFR2b signaling pathway, wherein a mesenchymal FGF10 gradient acts as a chemoattractant for the endoderm and promotes its proliferation (Bellusci et al., 1997; Park et al., 1998). *Fgf10* expression is strictly controlled by endodermal SHH pathways that function to finely tune *Fgf10* expression, assuring the correct size and shape of the growing buds (Pepicelli et al., 1998; Warburton, 2008). The growth of the developing lung must be continuously calibrated. In the mesenchyme, for example, fibronectin deposition is strictly controlled by WNT signaling during the early

stages of branching morphogenesis, and alterations of such regulation lead to impaired branching with enlarged terminal buds (De Langhe et al., 2005).

Cadherins are a large family of glycoproteins that mediate specific cell–cell adhesion in a Ca²⁺-dependent manner (Van Roy F. and Berx G., 2008). E-Cadherin (CDH1) is an important cadherin that is highly expressed in the epithelium throughout lung development. Structurally, CDH1 is comprised of a single transmembrane domain, an ectodomain for binding to other adjacent cells (Leckband and Prakasam, 2006), and a cytoplasmic domain that interacts with a variety of molecules that link CDH1 to the actin cytoskeleton, and hence to cell signaling and trafficking mechanisms (Bryant and Stow, 2004; Mege et al., 2006; Yap and Kovacs, 2003). *Cdh1* expression is promoted by transcription factors that bind cis-regulatory elements present in introns 1 and 2 (Goomer et al., 1994; Hennig et al., 1996; Liu et al., 2005), and can also be downregulated by specific transcription factors and corepressors (Peinado et al., 2007; Shi et al., 2003). CDH1 can undergo endocytosis and recycling through an endosomal

* Corresponding author. Fax: +1 323 361 3613.

E-mail address: dwarburton@chla.usc.edu (D. Warburton).

pathway that modulates its expression at the cell surface during dynamic morphogenetic processes (Bryant and Stow, 2004). CDH1 endocytosis, ubiquitination, and transport to the cell surface are finely regulated, adding another level of control over its activity (Davis et al., 2003; Palacios et al., 2002; Palacios et al., 2005). The regulation of *Cdh1* and its network of regulatory molecules has been widely investigated at the transcriptional and post-translational levels, but the effect of translational regulation is poorly understood.

An emerging understanding of the functional roles of microRNAs (miRNAs) in the post-transcriptional regulation of gene expression has shed light on the growing importance of these sub-cellular interactions. MiRNAs are small non-coding RNAs that can bind the 3'-UTR of target mRNAs and thus specifically inhibit their translation. The function of miRNAs can vary from acting as a molecular switch for specific events, to finely tuning or maintaining the stability and robustness of specific states. MiRNAs can also function as a control to eliminate specific gene expression on a translational level in specific locations (Brennecke et al., 2003; Cohen et al., 2006; Hornstein et al., 2005; Johnston and Hobert, 2003; Kwon et al., 2005; Sokol and Ambros, 2005).

In the lung, tissue-specific deletion of Dicer, the enzyme responsible for producing active miRNAs, results in branching arrest, revealing the importance of miRNAs for epithelial morphogenesis (Harris et al., 2006). The miR-17 family of miRNA, is comprised of three paralog clusters (miR-17–92, miR-106a–363, and miR-106b–25). The miR17–92 cluster has recently been described for its importance during lung development. In mice the miR-17–92 cluster is located on chromosome 14 and contains six miRNAs (miR-17, miR-18a, miR-19a, miR-20a, miR-19b-1, and miR-92–1). The miR-106a–363 cluster is located on mouse chromosome X and contains six miRNAs (miR-106a, miR-18b, miR-20b, miR-19b-2, miR-92–2, and miR-363), while miR-106b–25 is located on mouse chromosome 5 and contains three miRNAs (miR-106b, miR-93, and miR-25). Transgenic over-expression of the miR-17–92 cluster has been produced, showing its importance in controlling differentiation and proliferation of lung progenitor cells (Lu et al., 2007). Targeted deletion of the miR-17–92 cluster has been shown to produce smaller embryos with post-natal lethality, characterized by severely hypoplastic lungs. Furthermore during combined deletion of miR-17–92 cluster and its paralog miR-106b–25, mice showed earlier lethality before embryonic day 15 (Ventura et al., 2008).

Herein we report a specific functional role for miR-17, miR-20a, and miR-106b, in maintaining the structural homeostasis of developing lung epithelium. We show that miR-17, miR-20a, and miR-106b, target *Mapk14* and *Stat3* and thereby contribute to correctly modulating CDH1 expression in the epithelium in response to FGF10–FGFR2b signaling.

Materials and methods

RT-PCR and Real-time PCR

Total RNA was extracted using a miRNesasy Mini Kit (Qiagen) following the manufacturer's recommendations. For RNA extraction from lung tissue a pool of three samples was used. RT-PCR for mRNA was carried out using SuperScript II reverse transcriptase (Invitrogen) with random primers, while in RT-PCR for miRNA the TaqMan MicroRNA Reverse Transcription Kit (Applied Biosystems) was used. In both cases, reactions were assembled following the manufacturer's recommendations.

Real-time PCR was performed on a LightCycler 480 system (Roche). The universal probe library (Roche) was used for the analysis of mRNA expression. The miRNA Early Access Kit or Taqman microRNA assays (Applied Biosystems) were used for the screening of miRNAs differentially expressed during different stages of lung development. Primers and probes were used at a final concentration of 250 nM and

125 nM respectively. Following denaturation for 10 min at 95 °C, 45 cycles were performed with 10 s at 95 °C and 30 s at 60 °C. A relative quantification was performed using LightCycler 480 Software, and data were normalized with Actb for mRNA and U6 RNA for miRNA.

Immunoblotting

Samples were lysed in RIPA buffer (50 mM Tris–HCl, pH 7.4, 150 mM NaCl, 1 mM EDTA, 1% NP-40, 0.5% sodium deoxycholate, and 0.1% SDS) containing 1:100 dilution of protease inhibitor cocktail set III (Calbiochem). Immunoblotting was performed as previously described (Ding et al., 2007). The following primary antibodies were used: CDH1 (BD Biosciences) was used at 1:1000, STAT3 (Cell Signaling) was used at 1:2000, and MAPK14 (Cell Signaling) was used at 1:2000. ACTB (Chemicon) was used as a loading control and was used at 1:2000. Protein quantification was produced by densitometry analysis with the software Image J (Rasband W.S., U. S. National Institutes of Health, Bethesda, Maryland, USA, <http://rsb.info.nih.gov/ij/>).

In situ hybridization

Embryonic lung and epithelial lung explants were fixed in 4% paraformaldehyde (PFA) in phosphate-buffered saline (PBS) with rocking at 4 °C for 2 h and 5 min respectively, washed twice for 5 min in PBS at 4 °C, dehydrated in several washes with 70% ethanol (EtOH), and stored in absolute EtOH at –20 °C. For the in situ hybridization of miR-17 we used a modified protocol (Winnier et al., 1995) after taking the short size of mature miRNA into consideration. Samples were washed in Tris-Buffered Saline (TBS)/Tween-20 0.1% (TBST), digested with Proteinase K, fixed again in 4% PFA and washed in TBST. The samples were then transferred to hybridization buffer (Formamide 50%, SSC 5×, tRNA 0.05 mg/ml, sodium dodecyl sulfate 1%, Heparin 0.05 mg/ml) for 1 h at 50 °C and then hybridized with the probe at 55 °C over night. Locked Nucleic Acid (LNA) probes (Exiqon) were used to increase the stability of the hybridization. The next day the lung was washed with standard washing solutions. All washes were performed at 50 °C. After the washes the lungs were added to blocking solution made with sheep serum for 1 h and then incubated over night at 4 °C with anti-DIG antibody. The color reaction was performed with BM Purple solution after washes with TBST and NTMT (NaCl 100 mM, Tris pH9.5 100 mM, Tween-20 0.1%).

Isolated epithelial explant culture

Lungs from E11.5 embryos were treated with dispase (BD Biosciences) for 5 min at 4 °C and then transferred to Dulbecco's Modified Eagle's Medium: Nutrient Mixture F-12 (DMEM:F12) with 10% fetal bovine serum (FBS) (Invitrogen) to block the enzymatic reaction. The epithelial buds were separated from mesenchyme using tungsten needles (Fine Science Tools), embedded in 200 µl of Matrigel (BD Biosciences) diluted 1:1 with DMEM:F12 serum-free media containing 250 ng/ml FGF10 (R and D Systems), and left to polymerize by incubation at 37 °C for 20 min. The same media used for the matrigel mix was then added to the surface of the polymerized gel, and the samples were incubated at 37 °C (modified from Bellusci et al. (1997)). Locked nucleic acid against miR-17 (LNA17) or scrambled (SCRA) (Exiqon), or miR-17 mimic molecules (MIM17) (Thermo Scientific), were added at 1.5 µM to the isolated epithelium in presence of 0.3% lipofectamine RNAiMAX (Invitrogen).

Image analysis

Phase-contrast images of the samples were recorded using a digital camera (Diagnostic Instruments) connected to a reversed phase-contrast microscope (Leica). The public domain software ImageJ was

broadened by routines specifically developed by us and used to process and analyze the images. The main processing steps applied to the images are shown in Fig. 2D. Briefly, after correction for uneven illumination, the image underwent grey level threshold and contour smoothing to obtain the binary image of the tissue area. By using binary thinning procedures (Seul et al., 2000), the skeleton of this image was also derived and pruned to remove small artifactual branches. The number of terminal points and the number of branching points of the binary skeleton were then estimated. The former corresponds to the number of recognizable tissue branches; the latter provides an index of structural complexity of the sample (Guidolin et al., 2004). Finally, by tracing a poly-line with vertices located at the middle-point of each terminal branch of the binary skeleton, the area of tissue forming branches was measured and expressed as a percent of the total tissue area.

The number of formed branches, percentage of tissue area involved in branching, and branching complexity (i.e. the number of branching points of the binary skeleton) were the morphometric parameters estimated from each of 15 samples per group. Data were processed by using statistical analysis software (GraphPad Prism 3.03, GraphPad Software Inc., San Diego, USA). $P \leq 0.05$ was considered statistically significant. Comparisons of morphometric parameters between LNA17 and SCRA-treated tissue cultures were performed by Student's two-sample *t*-test.

Cell culture

Mouse lung epithelial cells (MLE12) (ATCC) were grown in DMEM/F12 (Invitrogen) in the presence of transferrin at 5 µg/ml, hydrocortisone at 10 µM, β -estradiol at 10 µM, HEPES at 10 mM, and L-glutamine 2 mM (SIGMA). Cells were grown in presence of penicillin/streptomycin and 2% FBS. Mouse mammary gland epithelial cells (NMuMG) (ATCC) were grown in DMEM in the presence of glucose at 4.5 g/l, sodium bicarbonate at 1.5 g/l, bovine insulin at 0.01 mg/ml (SIGMA) and 10% FBS.

Cell transfection and transduction

Modulation of miR-17 paralogs was obtained by use of LNA17 or SCRA, or MIM17. For *Stat3* or *Mapk14* silencing pools of specifically optimized small interfering RNA (si*Mapk14* and si*Stat3*) and scrambled siRNA (si*Scra*) were used (On-Target plus siRNA, Thermo Scientific). Molecules were added at 5 nM to the cells in presence of 0.1% lipofectamine RNAiMAX (Invitrogen) in media without antibiotics. After 5 h media was replaced with fresh complete media and cells grown for 36 h. Cells were then starved for 12 h with minimum media without serum in presence of 250 ng/ml FGF10 and then harvested for the analysis or fixed for staining. Adenoviruses for *Mapk14*, *Stat3* or GFP were added 24 h after plating of the cells at 50% confluence and then the same procedure used after transfection was followed. Experiments were repeated in triplicate. Adenoviruses having comparable titers were used. Effectiveness of siRNAs and adenoviruses were tested by real-time PCR on MLE12 cells.

Immunohistochemistry for whole-mount and sectioned tissue

Epithelial lung explants were fixed in 4% PFA, and hydrated by washing several times in Wash PBS (PBS containing 0.05% Tween-20, 0.1% BSA and 0.02% Sodium Azide). Tissues were then pre-incubated with 0.1% Triton X-100 in PBS for 5 min for membrane permeabilization, and washed 5× with Wash PBS. Tissues were then incubated in FBS diluted 1/20 in PBS for 45 min. Then, tissues were drained and incubated overnight at 4 °C with CDH1 antibody at 1:50 diluted in Wash PBS. The following day, samples were rinsed 5 times in Wash PBS. For the analysis on slides, tissues were fixed in 4% PFA, gradually dehydrated in ethanol, impregnated with toluene, embedded in

paraffin, and sectioned into 5 µm slices on poly-L-lysine coated slides. Antigen retrieval was performed by boiling the sample for 12 min in Na-citrate buffer (10 mM, pH 6.0) (Vector). CDH1 antibody (Ab) (BD Biosciences) was used at 1:100, and MAPK14 antibody (Cell Signaling) was used at 1:20. Incubation at 4 °C was performed in TBST with 3% bovine serum albumin. Secondary fluorescence antibodies (Jackson ImmunoResearch) were used for protein detection. DAPI (a nuclear dye of 4,6-diamidino-2-phenylindole; Vector Labs) was used to counterstain the DNA. Photomicrographs were taken using a Leica DMRA fluorescence microscope with a Hamamatsu digital camera. Digital analysis of CDH1 intensity and distribution was performed with the software MetaMorph (Molecular Devices). Cells analyzed in four different epithelial lung explants for each condition were analyzed. Data were reported as means \pm SEM and $P \leq 0.05$ was considered statistically significant.

β -galactosidase staining

TOPGAL mice were obtained from The Jackson Laboratory. TOPGAL epithelial explants were fixed in 4% paraformaldehyde (PFA) in phosphate-buffered saline (PBS) at 4 °C for 5 min with rocking, washed twice for 5 min in PBS at 4 °C, then transferred into freshly prepared X-gal solution and stained at 37 °C until a clear precipitate formed, modified from (Hogan et al., 1994). After rinsing with PBS, epithelial explants were post-fixed in 4% PFA in PBS. For vibratome sections, samples were embedded in a mixture of 300 mg/ml albumin, 5 mg/ml gelatin, and 0.6% glutaraldehyde, and sectioned at 30 µm.

Proliferation analysis

Ki67 antibody (Labvision) was used at 1:200, and signal was developed with the Histostatin Rabbit Primary Kit (Zymed-Invitrogen) following the manufacturer's instructions. Slides were counterstained with Hematoxylin. The total number of cells and the number of Ki67 positive cells in the epithelial lung explants were scored in 6 photomicrographs at the tip portions of 3 different sections in each of 3 MIM17-, 3 LNA17- and 3 SCRA-treated epithelial lung explants. The significance of the difference in proliferation between SCRA-treated and MIM17- or LNA17-treated samples was evaluated by Student's two-sample *t*-test.

Migration assay

A colorimetric chemotaxis cell migration assay (Millipore) was used for this analysis according with the manufacturer protocol. Briefly, MLE12 cells were transfected with SCRA, LNA17, and MIM17. After 48 h cells were harvested and plated at 30% confluence into 8 µm pore size membrane and grown in DMEM:F12 without FBS. After 24 h fresh media containing 250 ng/ml FGF10 was added, and after 14 h cells were stained and spectrophotometer reading at 560 nm (ND-1000, Nanodrop) was performed. Three experiments were performed for each group.

Adenoviral constructs

Adenoviral constructs for *Stat3* and *Mapk14* were produced using an adenoviral vector carrying a CMV promoter (ViraPower Adenoviral Expression System, Invitrogen). Coding sequences were NM_011951 for *Mapk14* and NM_213659 for *Stat3* and were cloned into an entry vector (pENTR-11, Invitrogen) before being transferred to the final vector. Adenoviruses were produced with HEK293A cells carrying E1a and E1b proteins necessary to generate the adenovirus. After repeated cycles of amplification, the adenoviruses were concentrated using a spin column based system (Cell Biolabs) and titered using an immunoassay with an antibody against hexon (Cell Biolabs). The titer obtained was between 10^7 and 10^8 transducing U/ml. Adenovirus




Fig. 1. MicroRNA expression profiles in embryonic versus adult lung and miR-17, miR-20a, and miR-106b spatial and temporal expression in embryonic lung. (A) Real-time PCR was done using primers that can identify 80 individual miRNAs, as listed in the panel, on total RNA extracted from embryonic (E10.5 and E11.5) versus adult lung to identify differentially expressed miRNAs. Results are shown as a heat map wherein green and red represent higher expression in the adult and embryonic lung respectively. miR-17 was clearly the most highly expressed in the early embryonic lung as compared to that of the adult. (B) Real-time PCR specifically tracking miR-17 in the embryonic, post-natal, and adult lung was carried out to determine its temporal expression. (C) Real-time PCR of miR-17 in the epithelium versus the mesenchyme of E12.5 embryonic lung was performed to determine its spatial localization within the embryonic lung (D–F), and the results were confirmed by in situ hybridization (ISH) in E12.5 embryonic lung. Whole-mount ISH (D, 3.2 and 14 \times) was further analyzed with vibratome sectioning (E, 3.2 and 14 \times). Control ISH was carried out using a probe composed of a scrambled sequence (F, 14 \times). (G) Real-time PCR specifically comparing miR-20a and miR-106b expression in the embryonic, post-natal, and adult lung is shown. (H) Real-time PCR of miR-20a and miR-106b in the epithelium versus the mesenchyme of E12.5 embryonic lung showing their relative spatial localization within the embryonic lung. Data in (B), (C), (G), and (H) are reported as means \pm SEM. * = $P \leq 0.05$. mes, mesenchyme; epi, epithelium; SCRA, scrambled probe.

containing GFP was used as a control. Experiments were repeated in triplicate with adenoviruses having comparable titers.

Computational target prediction

Stat3 and *Mapk14* were identified as principal miR17 targets based on the on-line available prediction databases TargetScan (<http://www.targetscan.org/>) and Miranda (<http://www.microrna.org/microrna/>). Seventy base pairs at the 5' UTR and 3' UTR of the miR-

17 binding position were further analyzed with the software mfold (Mathews et al., 1999; Zuker, 2003) (<http://mfold.bioinfo.rpi.edu/cgi-bin/rna-form1.cgi>) to determine the secondary structure and free energy. As a threshold for mouse sequences we used the previously reported value $dG = -13.4 \text{ kcal mol}^{-1}$ (Zhao et al., 2005). Stems $\geq 10 \text{ bp}$ were considered stabilizing elements, while hairpin loops $\geq 11 \text{ bp}$, interior loops $\geq 9 \text{ bp}$, bulge loops $\geq 7 \text{ bp}$, multiple branching loops $\geq 11 \text{ bp}$, and joint sequences and free ends $\geq 11 \text{ bp}$ were considered destabilizing elements (Zhao et al., 2005).




Fig. 2. miR-17, miR-20a, and miR-106b downregulation disrupted progression of embryonic lung epithelium branching. (A) Real-time PCR using total RNA extracted from isolated epithelial explants grown for 48 h in the presence of locked nucleic acid, scrambled (SCRA) or miR-17 specific (LNA17) was performed to determine the specificity of the LNA17 knockout probe. miR-17, miR-20a and miR-106b (three paralogs of the miR-17 family with high sequence conservation) showed significantly reduced expression after LNA17 administration, while miR-19b, a control member of the miR-17–92 cluster, was not affected. Data are reported as means \pm SEM. (B, 40 \times) Fluorescein-labeled LNA17 probe was transfected into the epithelial lung explants and was visible after 36 h. (C, 12 \times and 4 \times) Epithelial lung explants were grown for 72 h in a matrigel-based three-dimensional scaffold in the presence of 250 ng/ml FGF10. SCRA administration did not alter the normal pattern of branching progression (left two images), while LNA17 interfered with branching in the presence of FGF10 (right two images). (D) Morphometric parameters of LNA17- and SCRA-treated epithelial lung explants were collected and analyzed to determine their statistical significance. To the left, a description of the processing steps is depicted: the image of the tissue was corrected for uneven illumination and thresholded to obtain a binary image of the tissue area. The binary skeleton of this image was estimated and terminal branches (yellow) and branching points (blue) were identified. They are shown superimposed onto the image of the tissue area. The fraction of tissue involved in branching was then objectively defined as the area external to the poly-line with vertices located at the midpoint of each terminal branch of the binary skeleton (red). To the right, the morphometry from fifteen samples per group was analyzed. Data are reported as means \pm SEM. * = $P \leq 0.05$.





Fig. 3. CDH1 distribution is disrupted after miR-17, miR-20a, and miR-106b downregulation. (A, 20 \times and 60 \times) NMuMG cells were transfected with LNA17, SCRA, or LNA17 and MIM17. CDH1 IHC was performed to determine the impact of miR-17 downregulation on CDH1 distribution (green). Nuclei were stained with DAPI (blue). (B, 63 \times) Whole-mount IHC was performed using epithelial lung explants transfected with LNA17, SCRA, or LNA17 and MIM17, and grown for 72 h in presence of 250 ng/ml FGF10 to monitor changes in CDH1 distribution. (C) Real-time PCR using epithelial lung explants from the treatment described in (B) was carried out to determine variations in *Cdh1* mRNA levels. (D) Immunoblotting analysis was performed using whole-cell extract from NMuMG cells treated as described in (A) to monitor *Cdh1* changes at the translational level. (E, 63 \times and 330 \times) IHC was performed for CDH1 (red) using 5 μ m sections from epithelial lung explants treated as described in (B). On the left, images of terminal buds from scrambled LNA (top) or LNA17 (bottom) treated epithelial explants. On the right, electronic magnification of single cells showing the selected area (dashed lines) used to analyze CDH1 signal intensity and distribution (F, top). Data from cells analyzed in four different epithelial lung explants for each condition were analyzed (F, bottom). Data are reported as means \pm SEM. * = $P \leq 0.05$.






Fig. 5. miR-17 paralogs directly targets the 3'-UTR of *Stat3* and *Mapk14*. (A) *Stat3* and *Mapk14* temporal differential expression was determined by real-time PCR performed on total RNA extracted from embryonic, post-natal, and adult lungs. (B) miR-17, miR-20a, and miR-106b binding to the 3' UTR of *Stat3* and *Mapk14* are shown. (C) 70 bp miR-17 binding position flanking regions secondary structure and free energy (dG) are shown for *Stat3* and *Mapk14* based on mfold analysis. (D) MLE12 cells were transfected with expression vectors containing the luciferase open reading frame coupled with one, two, or three copies of the mutated (MU) or wild type (WT) miR-17 paralogs binding sequence of *Stat3* or *Mapk14*. Co-transfection with *Renilla* luciferase-expressing vector was used as a system to normalize luminescence levels. Data are reported as means \pm SEM. * = $P \leq 0.05$. (E) Immunoblotting from MLE12 cells whole extract after treatment with SCRA or LNA17 was performed to determine miR-17 paralogs translational regulation of *Stat3* and *Mapk14*.

epithelial branching is disrupted rather than the initiation of individual early branching events (Fig. 2D).

Altered expression and distribution of CDH1 after miR-17, miR-20a and miR-106b downregulation

In order to determine whether the observed phenotype was related to any alteration in epithelial cell-cell adhesion, we analyzed CDH1 expression. Epithelial cells transfected with LNA17 showed a

sharp loss of CDH1 distribution at their interacting edges compared with SCRA transfected epithelial cells. The resulting cell shape was rounded with diffuse CDH1 expression within the cell. In experiments in which MIM17 was co-transfected with LNA17, a rescue of the phenotype was observed (Fig. 3A). Similar results and rescue were obtained with epithelial lung explants after transfection with LNA17 alone or co-transfection of LNA17 with MIM17 (Fig. 3B). *Cdh1* changes at transcriptional level were not observed after LNA17 treatment (Fig. 3C), while differences of protein levels were observed after both

Table 1

Analysis of predicted miR-17, miR-20a, and miR-106b binding sites flanking UTRs of *Mapk14* and *Stat3*.

3'-UTR sequence	mFold dG (kcal/mol)		Structure	
	5' 70 bp	3' 70 bp	SE	DSE
<i>Mapk14</i>	−9.9	−11.4	–	HL, FE, MBL
<i>Stat3</i>	−5.4	−8.9	–	HL, J

Free energy (dG) of the 5'-UTR and 3' UTR flanking the miR-17 predicted binding position for *Mapk14* and *Stat3*. No Stabilizing Elements (SE) were present. Destabilizing Elements (DSE) are indicated: HL, Hairpin Loop; FE, Free End; MBL, multi-branching loop; J, Joint.

LNA17 and MIM17 (Fig. 3D). Analysis of sections from epithelial lung explants confirmed these alterations in epithelial cell shape and CDH1 distribution (Fig. 3E). Quantitative analysis on single cells showed that LNA17 treatment was characterized by lower expression and

increased lateral diffusion of CDH1 at the cell edges, the points of cell-to-cell interaction (Fig. 3F).

Downstream effects of miR-17, miR-20a and miR-106b downregulation on epithelial beta-catenin activity, migration, and proliferation

We used epithelial explants obtained from TOPGAL transgenic mice to determine if the alteration of Cdh1 expression observed after miR-17, miR-20a, and miR-106b downregulation affected beta-catenin activity. Transfection with LNA17 produced increased beta-catenin activity as shown by augmented TOPGAL expression in the branching epithelium. This phenotype was rescued after co-transfection of LNA17 and MIM17 (Fig. 4A). Furthermore, direct beta-catenin targets such as *Bmp4* and *Fgfr2b* were positively regulated after miR-17, miR-20a, and miR-106b downregulation, supporting the functional importance of these changes in beta-catenin activity (Fig. 4B). *Bmp4* overexpression in embryonic lung epithelium driven by SPC promoter has been reported and resulted




Fig. 6. Simultaneous modulation of *Stat3* and *Mapk14* mimics miR-17 paralogs effect. (A) Real-time PCR was performed on total RNA extracted from MLE12 cells after transduction with *Stat3* and *Mapk14* adenovirus to determine the effectiveness of the adenoviral constructs. Data were compared to results from MLE12 cells transduced with a control adenovirus. Data are reported as means \pm SEM. * = $P \leq 0.05$. (B, 20 \times) IHC for CDH1 (red) and MAPK14 (green) was performed in MLE12 cells after the overexpression of *Stat3* and *Mapk14* by transduction with their specific adenovirus. (C) Immunoblotting from MLE12 cells' whole extract after inhibition of *Stat3* and *Mapk14* by siRNA transfection was used to determine changes in CDH1.

in diminished branching with dilated terminal buds (Bellusci et al., 1996). Further, in vitro administration of BMP4 to isolated endoderm migrating toward FGF10-loaded beads has been shown to inhibit bud outgrowth (Weaver et al., 2000). Our observed increase in the area of the branches and disruption of budding shape may be partially produced by the increase in *Bmp4*, while the increase in the number of secondary branches must be related to different regulators that we speculate may be in turn related to CDH1 modulation.

Since CDH1 plays a key role in controlling the ability of epithelial cells to dissociate and migrate, we examined changes in the migratory capability of epithelial cells after miR-17 alteration. Treatment of MLE12 cells with LNA17 showed increased (60%) migration during FGF10 stimulation, compared with cells treated with SCRA. Meanwhile, co-transfection with both LNA17 and MIM17 completely rescued the ability of MLE12 cells to migrate towards FGF10, while transfection with MIM17 alone did not produce any appreciable effect (Fig. 4D).

The branching alteration observed during LNA17 treatment may depend on changes in epithelial cell proliferation. To address this possibility, we analyzed cell proliferation in epithelial lung explants after LNA17 treatment as compared to proliferation after MIM17 administration. Treatment with LNA17 did not show significant changes in epithelial proliferation compared with samples treated with SCRA. However MIM17 transfection produced a significant (40%) increase of epithelial proliferation (Fig. 4C). Increased cellular proliferation was previously reported in transgenic mice misexpressing the entire miR-17–92 cluster (Lu et al., 2007). Both MIM17 and LNA17 treated samples, analyzed with TUNEL assay, did not show signs of apoptosis (data not shown).

Mapk14 and Stat3 are direct targets of miR-17 paralogs at the translational level, and their simultaneous overexpression mimics miR-17 paralogs downregulation

In order to identify transcripts directly regulated by miR-17 paralogs we analyzed available miRNA target prediction software for putative miR-17, miR-20a and miR-106b target genes (TargetScan and Miranda). Candidate genes were analyzed at the target position flanking regions to determine their level of accessibility with the software mfold. Two genes, *Mapk14* and *Stat3*, were shown to possess a free energy higher than the characteristic threshold level for mouse models ($\Delta G = -13.4 \text{ kcal mol}^{-1}$, Zhao et al. 2005) (Fig. 5C) and to have destabilizing elements in their secondary structure that may facilitate the accessibility of miR-17, miR-20a, and miR-106b to the binding site (Table 1). The levels of expression of both *Stat3* and *Mapk14* in embryonic lung development are stable during the pseudoglandular stage (Fig. 5A), suggesting that their stabilization may be important for correct branching morphogenesis. To determine the ability of miR-17 paralogs to directly bind to the 3'-UTR of these two genes (Fig. 5B), we produced luciferase-expressing vectors containing respectively one, two, or three copies of their target position. Vectors containing one, two or three copies of the target position mutated inside the "seed" sequence were used as reference for the quantification of luciferase expression. Transfection of the luciferase vectors into MLE12 epithelial cells showed a progressive reduction in luminescence. For *Stat3* in the presence of one, two, or three copies of the target sequence, 29%, 32%, or 50% luminescence reduction was observed, while for *Mapk14* the decrease was respectively of 20%, 37% or 46%. This result demonstrates the specificity of the binding of miR-17, miR-20a and miR-106b to the 3'-UTR of *Stat3* and *Mapk14* (Fig. 5D).

Regulation of *Stat3* and *Mapk14* expression at the translational level was then analyzed by western blotting after miR-17, miR-20a, and miR-106b downregulation, confirming the expected changes in protein levels (Fig. 5E).

To determine the ability of *Stat3* and *Mapk14* to control *Cdh1* expression, we used adenoviral overexpression approaches in MLE12 epithelial cells. The effectiveness of the adenoviral construct was determined by real-time PCR (Fig. 6A). While the individual overexpression of *Stat3* or *Mapk14* did not produce evident changes in CDH1 distribution (data not shown), co-overexpression of both molecules strongly replicated the effect observed on CDH1 after LNA17 administration, and co-staining of epithelial cells with CDH1 and MAPK14 showed an inverse correlation between CDH1 levels and MAPK14 expression (Fig. 6B). Furthermore while downregulation of *Stat3* alone (data not shown) or *Mapk14* alone (Fig. 6C) with specific siRNAs did not modify CDH1 expression, downregulation of both *Stat3* and *Mapk14* showed specific overexpression of CDH1 (Fig. 6C).

We therefore conclude that miR-17 family of miRNA, specifically miR-17, miR-20a and miR-106b, negatively modulate FGF10-FGFR2b downstream signaling by specifically targeting *Stat3* and *Mapk14*, hence regulating activation of *Cdh1* and beta-catenin, which in turn modulate epithelial bud morphogenesis in response to FGF10 signaling.

Discussion

In this report we have identified a specific role of miR-17 family of miRNAs in maintaining homeostasis of epithelial structure within embryonic lung during branching morphogenesis. We used in vitro gain and loss of function approaches to test our hypothesis that an important function of miR-17, miR-20a, and miR-106b is to maintain correct CDH1 distribution during branching morphogenesis. Changes in branching and epithelial homeostasis were found during miR-17 paralogs loss of function experiments and these were associated with changes in *Cdh1* expression. Furthermore, we determined that miR-17, miR-20a, and miR-106b directly target both *Mapk14* and *Stat3*, which in turn regulate CDH1.

Dynamic adhesive mechanisms are important for the maintenance of correct epithelial tissue architecture during development. Cell–cell interaction in the epithelium of the developing lung is subject to dynamic changes during growth and specification of branching patterns (Warburton, 2008). CDH1 is a key adhesion receptor for epithelial cell identification and grouping during development; in its absence, the interaction between extracellular matrix, cytoplasmic plaques, and other adhesion molecules cannot be maintained. Mouse embryonic lung explants grown in the presence of a CDH1 antibody show an altered pattern of branching (Hirai et al., 1989). The expression of *Cdh1*, like other cadherins, is regulated at both transcriptional and post-translational levels and is controlled during trafficking and arrangement at the cell surface (Halbleib and Nelson, 2006). Therefore, precise control of *Cdh1* expression is necessary for cell–cell interaction during lung development. Herein, we provide evidence that miR-17, miR-20a, and miR-106b have a critical function to control *Mapk14* and *Stat3*, thus maintaining correct CDH1 levels and hence avoiding excessive loss of CDH1 during lung branching morphogenesis.

miR-17 and miR-20a expression is higher early in embryonic lung development and decreases with gestational age, miR-106b expression is also high during embryonic stages between E-12.5 and E16.5 and then becomes even higher again in the adult lung. Most importantly these three miRNAs are clearly present during the pseudoglandular stage when the embryonic lung undergoes repetitive budding to develop its characteristic three-dimensional architecture. *Mapk14* and *Stat3* also both maintain stable expression during the pseudoglandular stage. Taken together with our observation of a synergistic function of *Mapk14* and *Stat3*, this suggests that the stable and correct expression of both *Mapk14* and *Stat3* are important in maintaining the correct level of CDH1 during the early and critical stages of lung branching morphogenesis.

Mapk14 has been implicated in regulation of developmental processes such as wing development in *Drosophila* (Adachi-Yamada

et al., 1999) and recently its importance in maintaining a correct level of CDH1 degradation has been demonstrated in mice: disruption of MAPK14-interacting protein, a specific MAPK14 activator, can cause defects in neural tube and eye formation as well as gastrulation due to impaired CDH1 degradation (Zohn et al., 2006). *Mapk14* also has a functional role in lung branching morphogenesis. Both chemical inhibitors and short hairpin RNAs against *Mapk14* produced increased levels of CDH1, leading to impaired branching (Liu et al., 2008). These data are consistent with our results on *Mapk14* overexpression following miR-17 downregulation with LNA, in which CDH1 expression was notably altered. On the other hand, overexpression of *Mapk14* alone did not affect CDH1 localization (data not shown). Our observations were nevertheless consistent with the data of Liu et al. in which *Mapk14* downregulation showed changes in *Cdh1* expression but not in its localization. Furthermore, our data on *Mapk14* and *Stat3* co-overexpression and co-downregulation reinforce the concept that both molecules are important in simultaneously controlling CDH1 distribution.

Stat3 has been implicated in the regulation of cell motility during reparative processes in the lung (Kida et al., 2008) and has an important role during branch formation in neural path finding (Conway, 2006). This correlates with our observation of increased migratory ability of MLE12 cells after miR-17 downregulation. Furthermore, *Stat3* has been shown to be essential during gastrulation movements in zebrafish because it is responsible for activating Snail, a negative regulator of *Cdh1* expression (Yamashita et al., 2004).

The effect of miR-17, miR-20a, and miR-106b downregulation in our in vitro system can be considered a multiple-target effect. Our results show that co-overexpression of *Mapk14* and *Stat3* can mimic the effect of miR-17 downregulation. Therefore, *Mapk14* and *Stat3* are two main molecules that must apparently interact to produce this phenotype.

Other targets of miR-17 have also been identified, but none of them seem to be major candidates for participating in the regulation of epithelial structural homeostasis. E2f1 is an important regulator of apoptosis, but lung development is characterized by the relatively complete absence of apoptosis. In our system, no changes in apoptosis were observed after up or down regulation of miR-17 (data not shown). AIB1 is a proto-oncogene, and cells derived from mice in which this gene was deleted showed decreased proliferation in response to insulin-like growth factor 1 (Wang et al., 2000). Since miR-17 overexpression produced a hyperproliferative effect on epithelial cells, AIB1 cannot be logically implicated in our model. Rbl2 has been proposed as a direct target of miR-17 (Lu et al., 2007), and its deletion produce epithelial cell hyperplasia. Downregulation of Rbl2 in our experiments of miR-17 overexpression could conceivably contribute to the hyper-cellularity that we have observed. However Rbl2 expression becomes important in the lung only during late embryonic stages and is higher in the adult. Therefore, its major involvement during the early stages of lung development is unlikely.

Recently miR-17 has been suggested to have the ability to regulate the G1/S cell cycle checkpoint by targeting both promoters and inhibitors of proliferation in different contexts (Cloonan et al., 2008). While blocking the effects of proliferation inhibitors such as MAPK9 or p21 may account for increased proliferation after miR-17 overexpression, in our system inhibition of miR-17 did not produce an arrest of proliferation. This suggests that the changes in *Cdh1* expression and localization that we observed are not likely to be caused primarily by changes in cell cycle regulation.

In vivo overexpression of the miR-17–92 cluster has been reported to produce hyper-cellularity in the lung (Liu et al., 2007). These data are consistent with our results in which specific overexpression of miR-17 produced increased cell proliferation.

Hypoplastic lung at embryonic stage E18 was also noted in the presence of in vivo deletion of the entire miR-17–92 cluster.

Furthermore double knockout for miR-17–92 and miR-106b–25 clusters showed a more severe phenotype with lethality before embryonic day 15 (Ventura et al., 2008). In the single deletion of miR-17–92 cluster the authors state that even though hypoplastic lung was observed no branching defects were observed. However, no detailed analysis or discussion of embryonic lung phenotypes in the double knockout were done therein. Our observations on alteration of embryonic lung epithelium structural homeostasis after co-downregulation of miR-17, miR-20a and miR-106b, suggest that the double knock out of miR-17–92 and miR-106b–25 cluster may indeed have embryonic lung developmental abnormalities.

We therefore propose a model in which miR-17, miR-20a, and miR-106b functions to maintain the correct level of expression and distribution of CDH1 in embryonic lung epithelial cells through negative modulation of *Mapk14* and *Stat3*. Thus, if miR-17 paralogs are eliminated or inhibited, the structural integrity of the process of epithelial branching becomes compromised.

Conclusions

In summary we have identified a specific functional role of miR-17, miR-20a, and miR-106b to control the structural homeostasis of the epithelium during early embryonic branching morphogenesis in the lung. The dual targets that we have identified are *Mapk14* and *Stat3*, which can mimic the phenotype elicited by miR-17, miR-20a, and miR-106b co-inhibition, when they are overexpressed together. The detailed molecular links between these two targets and the observed alteration of *Cdh1* expression remain to be further defined.

Acknowledgments

We would like to thank Dr. Yannan Ouyang of cellular imaging core at the Saban Research Institute for the discussion of image analysis. HL60231, HL44977 and HL44060 to DW and American Heart Association Western Affiliates Senior Postdoctoral Fellowship to GC.

References

- Adachi-Yamada, T., Nakamura, M., Irie, K., Tomoyasu, Y., Sano, Y., Mori, E., Goto, S., Ueno, N., Nishida, Y., Matsumoto, K., 1999. p38 mitogen-activated protein kinase can be involved in transforming growth factor beta superfamily signal transduction in *Drosophila* wing morphogenesis. *Mol. Cell. Biol.* 19, 2322–2329.
- Akao, Y., Nakagawa, Y., Naoe, T., 2006. let-7 microRNA functions as a potential growth suppressor in human colon cancer cells. *Biol. Pharm. Bull.* 29, 903–906.
- Bellusci, S., Henderson, R., Winnier, G., Oikawa, T., Hogan, B.L., 1996. Evidence from normal expression and targeted misexpression that bone morphogenetic protein (Bmp-4) plays a role in mouse embryonic lung morphogenesis. *Development* 122, 1693–1702.
- Bellusci, S., Grindley, J., Emoto, H., Itoh, N., Hogan, B.L., 1997. Fibroblast growth factor 10 (FGF10) and branching morphogenesis in the embryonic mouse lung. *Development* 124, 4867–4878.
- Brennecke, J., Hipfner, D.R., Stark, A., Russell, R.B., Cohen, S.M., 2003. bantam encodes a developmentally regulated microRNA that controls cell proliferation and regulates the proapoptotic gene *hid* in *Drosophila*. *Cell* 113, 25–36.
- Bryant, D.M., Stow, J.L., 2004. The ins and outs of E-cadherin trafficking. *Trends Cell Biol.* 14, 427–434.
- Cardoso, W.V., Lu, J., 2006. Regulation of early lung morphogenesis: questions, facts and controversies. *Development* 133, 1611–1624.
- Cloonan, N., Brown, M.K., Steptoe, A.L., Wani, S., Chan, W.L., Forrest, A.R., Kolle, G., Gabrielli, B., Grimmond, S.M., 2008. The miR-17–5p microRNA is a key regulator of the G1/S phase cell cycle transition. *Genome Biol.* 9, R127.
- Cohen, S.M., Brennecke, J., Stark, A., 2006. Denoising feedback loops by thresholding—a new role for microRNAs. *Genes Dev.* 20, 2769–2772.
- Conway, G., 2006. STAT3-dependent pathfinding and control of axonal branching and target selection. *Dev. Biol.* 296, 119–136.
- Davis, M.A., Ireton, R.C., Reynolds, A.B., 2003. A core function for p120-catenin in cadherin turnover. *J. Cell. Biol.* 163, 525–534.
- De Langhe, S.P., Sala, F.G., Del Moral, P.M., Fairbanks, T.J., Yamada, K.M., Warburton, D., Burns, R.C., Bellusci, S., 2005. Dickkopf-1 (DKK1) reveals that fibronectin is a major target of Wnt signaling in branching morphogenesis of the mouse embryonic lung. *Dev. Biol.* 277, 316–331.
- Ding, W., Shi, W., Bellusci, S., Groffen, J., Heisterkamp, N., Minoo, P., Warburton, D., 2007. Sprout2 downregulation plays a pivotal role in mediating crosstalk between TGF-beta1 signaling and EGF as well as FGF receptor tyrosine kinase-ERK pathways in mesenchymal cells. *J. Cell. Physiol.* 212, 796–806.

- Fish, J.E., Santoro, M.M., Morton, S.U., Yu, S., Yeh, R.F., Wythe, J.D., Ivey, K.N., Bruneau, B.G., Stainier, D.Y., Srivastava, D., 2008. miR-126 regulates angiogenic signaling and vascular integrity. *Dev. Cell* 15, 272–284.
- Goomer, R.S., Holst, B.D., Wood, I.C., Jones, F.S., Edelman, G.M., 1994. Regulation in vitro of an L-CAM enhancer by homeobox genes HoxD9 and HNF-1. *Proc. Natl. Acad. Sci. U. S. A.* 91, 7985–7989.
- Guidolin, D., Vacca, A., Nussdorfer, G.G., Ribatti, D., 2004. A new image analysis method based on topological and fractal parameters to evaluate the angiostatic activity of docetaxel by using the Matrigel assay in vitro. *Microvasc. Res.* 67, 117–124.
- Halbleib, J.M., Nelson, W.J., 2006. Cadherins in development: cell adhesion, sorting, and tissue morphogenesis. *Genes Dev.* 20, 3199–3214.
- Harris, K.S., Zhang, Z., McManus, M.T., Harfe, B.D., Sun, X., 2006. Dicer function is essential for lung epithelium morphogenesis. *Proc. Natl. Acad. Sci. U. S. A.* 103, 2208–2213.
- Hennig, G., Lowrick, O., Birchmeier, W., Behrens, J., 1996. Mechanisms identified in the transcriptional control of epithelial gene expression. *J. Biol. Chem.* 271, 595–602.
- Hirai, Y., Nose, A., Kobayashi, S., Takeichi, M., 1989. Expression and role of E- and P-cadherin adhesion molecules in embryonic histogenesis. II. Skin morphogenesis. *Development* 105, 271–277.
- Hogan, B., Beddington, R., Constantini, F., Lacy, E., 1994. *Manipulating the Mouse Embryo*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Hornstein, E., Mansfield, J.H., Yekta, S., Hu, J.K., Harfe, B.D., McManus, M.T., Baskerville, S., Bartel, D.P., Tabin, C.J., 2005. The microRNA miR-196 acts upstream of Hoxb8 and Shh in limb development. *Nature* 438, 671–674.
- Johnston, R.J., Hobert, O., 2003. A microRNA controlling left/right neuronal asymmetry in *Caenorhabditis elegans*. *Nature* 426, 845–849.
- Kida, H., Mucenski, M.L., Thitoff, A.R., Le Cras, T.D., Park, K.S., Ikegami, M., Muller, W., Whitsett, J.A., 2008. GP130-STAT3 regulates epithelial cell migration and is required for repair of the bronchiolar epithelium. *Am. J. Pathol.* 172, 1542–1554.
- Kwon, C., Han, Z., Olson, E.N., Srivastava, D., 2005. MicroRNA1 influences cardiac differentiation in *Drosophila* and regulates Notch signaling. *Proc. Natl. Acad. Sci. U. S. A.* 102, 18986–18991.
- Leckband, D., Prakasam, A., 2006. Mechanism and dynamics of cadherin adhesion. *Annu. Rev. Biomed. Eng.* 8, 259–287.
- Liu, Y., Martinez, L., Ebine, K., Abe, M.K., 2008. Role for mitogen-activated protein kinase p38 alpha in lung epithelial branching morphogenesis. *Dev. Biol.* 314, 224–235.
- Liu, Y.N., Lee, W.W., Wang, C.Y., Chao, T.H., Chen, Y., Chen, J.H., 2005. Regulatory mechanisms controlling human E-cadherin gene expression. *Oncogene* 24, 8277–8290.
- Lu, Y., Thomson, J.M., Wong, H.Y., Hammond, S.M., Hogan, B.L., 2007. Transgenic overexpression of the microRNA miR-17-92 cluster promotes proliferation and inhibits differentiation of lung epithelial progenitor cells. *Dev. Biol.* 310, 442–453.
- Mathews, D.H., Sabina, J., Zuker, M., Turner, D.H., 1999. Expanded sequence dependence of thermodynamic parameters improves prediction of RNA secondary structure. *J. Mol. Biol.* 288, 911–940.
- Mege, R.M., Gavard, J., Lambert, M., 2006. Regulation of cell–cell junctions by the cytoskeleton. *Curr. Opin. Cell. Biol.* 18, 541–548.
- Palacios, F., Schweitzer, J.K., Boshans, R.L., D'Souza-Schorey, C., 2002. ARF6-GTP recruits Nm23-H1 to facilitate dynamin-mediated endocytosis during adherens junctions disassembly. *Nat. Cell Biol.* 4, 929–936.
- Palacios, F., Tushir, J.S., Fujita, Y., D'Souza-Schorey, C., 2005. Lysosomal targeting of E-cadherin: a unique mechanism for the down-regulation of cell–cell adhesion during epithelial to mesenchymal transitions. *Mol. Cell. Biol.* 25, 389–402.
- Park, W.Y., Miranda, B., Lebeche, D., Hashimoto, G., Cardoso, W.V., 1998. FGF-10 is a chemotactic factor for distal epithelial buds during lung development. *Dev. Biol.* 201, 125–134.
- Peinado, H., Olmeda, D., Cano, A., 2007. Snail, Zeb and bHLH factors in tumour progression: an alliance against the epithelial phenotype? *Nat. Rev. Cancer* 7, 415–428.
- Pepicelli, C.V., Lewis, P.M., McMahon, A.P., 1998. Sonic hedgehog regulates branching morphogenesis in the mammalian lung. *Curr. Biol.* 8, 1083–1086.
- Seul, M., O'Gorman, L., Sammon, M.J., 2000. *Practical Algorithms for Image Analysis*. Cambridge Univ. Press, New York.
- Shi, Y., Sawada, J., Sui, G., Affar el, B., Whetstone, J.R., Lan, F., Ogawa, H., Luke, M.P., Nakatani, Y., 2003. Coordinated histone modifications mediated by a CtBP co-repressor complex. *Nature* 422, 735–738.
- Sokol, N.S., Ambros, V., 2005. Mesodermally expressed *Drosophila* microRNA-1 is regulated by Twist and is required in muscles during larval growth. *Genes Dev.* 19, 2343–2354.
- Van Roy, F., Berx, G., 2008. The cell–cell adhesion molecule E-cadherin. *Cell. Mol. Life Sci.* 65, 3756–3788.
- Ventura, A., Young, A.G., Winslow, M.M., Lintault, L., Meissner, A., Erkland, S.J., Newman, J., Bronson, R.T., Crowley, D., Stone, J.R., Jaenisch, R., Sharp, P.A., Jacks, T., 2008. Targeted deletion reveals essential and overlapping functions of the miR-17 through 92 family of miRNA clusters. *Cell* 132, 875–886.
- Wang, Z., Rose, D.W., Hermanson, O., Liu, F., Herman, T., Wu, W., Sze, D., Gleiberman, A., Krones, A., Pratt, K., Rosenfeld, R., Glass, C.K., Rosenfeld, M.G., 2000. Regulation of somatic growth by the p160 coactivator p/CIP. *Proc. Natl. Acad. Sci. U. S. A.* 97, 13549–13554.
- Warburton, D., 2008. Developmental biology: order in the lung. *Nature* 453, 733–735.
- Warburton, D., Bellusci, S., De Langhe, S., Del Moral, P.M., Fleury, V., Mailleux, A., Tefft, D., Unbekandt, M., Wang, K., Shi, W., 2005. Molecular mechanisms of early lung specification and branching morphogenesis. *Pediatr. Res.* 57, 26R–37R.
- Weaver, M., Dunn, N.R., Hogan, B.L., 2000. Bmp4 and Fgf10 play opposing roles during lung bud morphogenesis. *Development* 127, 2695–2704.
- Winnier, G., Blessing, M., Labosky, P.A., Hogan, B.L., 1995. Bone morphogenetic protein-4 is required for mesoderm formation and patterning in the mouse. *Genes Dev.* 9, 2105–2116.
- Yamashita, S., Miyagi, C., Fukada, T., Kagara, N., Che, Y.S., Hirano, T., 2004. Zinc transporter LIV1 controls epithelial–mesenchymal transition in zebrafish gastrula organizer. *Nature* 20, 298–302.
- Yap, A.S., Kovacs, E.M., 2003. Direct cadherin-activated cell signaling: a view from the plasma membrane. *J. Cell. Biol.* 160, 11–16.
- Zhao, Y., Samal, E., Srivastava, D., 2005. Serum response factor regulates a muscle-specific microRNA that targets Hand2 during cardiogenesis. *Nature* 436, 214–220.
- Zohn, I.E., Li, Y., Skolnik, E.Y., Anderson, K.V., Han, J., Niswander, L., 2006. p38 and a p38-interacting protein are critical for downregulation of E-cadherin during mouse gastrulation. *Cell* 125, 957–969.
- Zuker, M., 2003. Mfold web server for nucleic acid folding and hybridization prediction. *Nucleic Acids Res.* 31, 3406–3415.